

Stimulus onset quenches neural variability: a widespread cortical phenomenon

Mark M Churchland^{1,2,16}, Byron M Yu^{1-3,16}, John P Cunningham¹, Leo P Sugrue^{2,4}, Marlene R Cohen^{2,4}, Greg S Corrado^{2,4}, William T Newsome^{2,4,5}, Andrew M Clark⁶, Paymon Hosseini⁶, Benjamin B Scott⁶, David C Bradley⁶, Matthew A Smith⁷, Adam Kohn^{8,9}, J Anthony Movshon⁹, Katherine M Armstrong^{2,5}, Tirin Moore^{2,5}, Steve W Chang¹⁰, Lawrence H Snyder¹⁰, Stephen G Lisberger¹¹, Nicholas J Priebe¹², Ian M Finn¹³, David Ferster¹³, Stephen I Ryu^{1,14}, Gopal Santhanam¹, Maneesh Sahani³ & Krishna V Shenoy^{1,2,15}

Neural responses are typically characterized by computing the mean firing rate, but response variability can exist across trials. Many studies have examined the effect of a stimulus on the mean response, but few have examined the effect on response variability. We measured neural variability in 13 extracellularly recorded datasets and one intracellularly recorded dataset from seven areas spanning the four cortical lobes in monkeys and cats. In every case, stimulus onset caused a decline in neural variability. This occurred even when the stimulus produced little change in mean firing rate. The variability decline was observed in membrane potential recordings, in the spiking of individual neurons and in correlated spiking variability measured with implanted 96-electrode arrays. The variability decline was observed for all stimuli tested, regardless of whether the animal was awake, behaving or anaesthetized. This widespread variability decline suggests a rather general property of cortex, that its state is stabilized by an input.

A fundamental approach of systems neuroscience is to probe the brain with repeated stimulus trials and infer neural mechanism from the recorded responses. Extracellularly recorded responses are typically analyzed by computing the average spike rate across trials. By averaging, the experimenter hopes to overcome the apparent noisiness of spiking and estimate the true change in the neuron's underlying firing rate. It is probably true that much of the recorded spiking variability is effectively noise and doesn't reflect fundamentally different responses on different trials. However, it is clear that the neural response can vary meaningfully across trials. For example, the neural state may be initially similar across trials but become variable in response to a stimulus¹. Alternately, sensory cortex can be restless and active² before stimulus onset. A central question is whether the stimulus-driven response suppresses such ongoing variability³⁻⁵, superimposes with it^{2,6,7} or yields even greater variability as a result of nonlinear interactions⁸? In general, does stimulus onset drive variability up (resulting from the variable responses themselves) or down (resulting from suppression of variable ongoing activity)?

The mean rate provides an incomplete characterization of the neural response. A fuller characterization requires, at the least, knowing whether rate variability is present and how it changes with time.

For example, fundamentally different response patterns can have similar across-trial means (Fig. 1a,b). The mean can also erroneously suggest that there is little stimulus-driven response (Fig. 1c, a similar scenario using a simulated network is shown in Supplementary Fig. 1). Because such situations may be common, it is important to characterize not only the stimulus-driven change in mean rate but also the stimulus-driven change in rate variance.

The effect of a stimulus on variability could depend on the brain area, stimulus and task. However, stimulus onset reduces both membrane potential variability in anaesthetized cat V1 (refs. 3,4) and firing-rate variability in premotor cortex of reaching monkeys^{9,10}. The presence of similar effects in two very different contexts suggests that a decline in variability could be a common feature of the cortical response. This would agree with recent theoretical work^{11,12} indicating that such an effect may be a general property of large recurrent networks.

To address this issue, we analyzed recordings from many cortical areas that were driven by a variety of stimuli. A measure of firing-rate variability (the Fano factor) revealed a stimulus-driven decline in variability that was similar in time course to the decline in V1 membrane-potential variability. This decline was present not only for anaesthetized V1 but for all of the cortical areas tested, regardless of the stimulus or

¹Department of Electrical Engineering, ²Neurosciences Program, Stanford University School of Medicine, Stanford University, Stanford, California, USA. ³Gatsby Computational Neuroscience Unit, University College London, London, UK. ⁴Howard Hughes Medical Institute, ⁵Department of Neurobiology, Stanford University School of Medicine, Stanford University, Stanford, California, USA. ⁶Department of Psychology and Committee on Computational Neuroscience, University of Chicago, Chicago, Illinois, USA. ⁷Department of Neuroscience and Center for the Neural Basis of Cognition, University of Pittsburgh, Pittsburgh, Pennsylvania, USA. ⁸Department of Neuroscience, Albert Einstein College of Medicine, Bronx, New York, USA. ⁹Center for Neural Science, New York University, New York, New York, USA. ¹⁰Department of Anatomy and Neurobiology, Washington University School of Medicine, St. Louis, Missouri, USA. ¹¹Howard Hughes Medical Institute, W.M. Keck Foundation Center for Integrative Neuroscience and Department of Physiology, University of California San Francisco, San Francisco, California, USA. ¹²Section of Neurobiology, School of Biological Sciences, University of Texas at Austin, Austin, Texas, USA. ¹³Department of Neurobiology and Physiology, Northwestern University, Evanston, Illinois, USA. ¹⁴Department of Neurosurgery, Stanford University School of Medicine, Stanford University, Stanford, California, USA. ¹⁵Department of Bioengineering, Stanford University, Stanford, California, USA. ¹⁶These authors contributed equally to this work. Correspondence should be addressed to M.M.C. (church@stanford.edu).

Received 21 September 2009; accepted 13 January 2010; published online 21 February 2010; doi:10.1038/nn.2501




Figure 1 Schematic illustration of possible types of across-trial firing rate variability. (a–c) We suppose that the same stimulus is delivered four times (four trials) yielding four different responses. a and b were constructed to have the same mean response across the four trials. Stimulus-driven decline in variability is shown in a. Stimulus-driven rise in variability is shown in b. Stimulus-driven decline in variability with little change in mean rate is shown in c.

behavioral state. The decline was also present in the correlated firing-rate variability of neurons recorded using implanted multi-electrode arrays. Finally, we determined how recently developed methods, when applied to simultaneous multi-electrode recordings, can reconstruct the variable evolution of firing rates on individual trials.

RESULTS

Across-trial variability in the membrane potential

Stimuli and task events can alter the structure and correlation¹³ of membrane-potential variability. In particular, visual stimuli drive a reduction in membrane potential (V_m) variability in cat primary visual cortex (V1) that is largely independent of stimulus orientation^{3,4}. We re-analyzed previously reported data⁴ to determine the time course of this effect (Fig. 2). Stimulus onset drives an immediate decline in V_m variability. This decline occurs even for nonpreferred stimuli that elicit little change in mean V_m (see also refs. 3,4). Average variability (across all neurons and conditions) declined rapidly following stimulus onset and then remained at a rough plateau (Fig. 2c). The variability in question was across-trial variability, with a fairly long autocorrelation. When V_m was low (or high), it tended to stay low (or high) for tens to hundreds of milliseconds.

The relationship between intracellularly recorded V_m variability and extracellularly recorded firing-rate variability is likely to be complex, given the nonlinear and dynamic relationship between

V_m and firing rate (for example, considerable V_m variability occurs below threshold). One nevertheless expects across-trial V_m variability to produce across-trial firing-rate variability. A larger question is whether the observed decline in variability is specific to V1 or whether it reflects a broader phenomenon. The latter is suggested by both the presence of a similar effect in premotor cortex^{9,10} and recent theoretical work^{11,12}.

Addressing these issue requires quantifying firing-rate variability in extracellular recordings. Although quantifying V_m variability is straightforward, quantifying firing-rate variability is more complicated. Extracellularly recorded spike trains are usually described in terms of an underlying firing rate (often termed λ) observed via a noisy point process (for example, Poisson) that produces spikes. It should be stressed that this conception captures the statistics of neurons embedded in a network^{14,15}; spike generation at the axon hillock is not responsible for the noisy spiking-process statistics¹⁶, nor is firing rate synonymous with membrane potential. Instead, the underlying firing rate can be thought of as the average response of many similarly tuned neurons or as the average response of one neuron across truly identical trials. Of course, repeated trials are not guaranteed to be truly identical; the underlying firing rate may differ somewhat. It is precisely this variability that we wished to capture, while ignoring variability arising from the roughly Poisson spiking. Spiking variability may have interesting structure of its own, but for present purposes, it acts as noise.

Poisson spiking-process noise can severely mask underlying firing-rate variability (Supplementary Fig. 1). It is therefore rarely possible to discern changes in firing-rate variability by eye. We used two approaches to isolate the underlying firing-rate variability from the variability contributed by spiking noise. The first approach employed a modified method for computing the Fano factor. This method is applicable to conventionally recorded single-neuron data, allowing analysis of a large number of existing datasets. The second approach used factor analysis to assess covariance in large-scale simultaneous recordings. These methods are technically very different, but both are intended to assess the same thing: the degree of across-trial firing rate variability, independent of the contribution of noisy spiking statistics.

A variability decline across multiple cortical areas

We first employed the Fano factor, which is the spike-count variance divided by the spike-count mean. Counts were made in a sliding





Figure 2 Analysis of intracellularly recorded membrane potential from cat V1. Stimuli were drifting sine-wave gratings presented at different orientations and frequencies. Spikes were removed before further analysis. Analysis employed a 50-ms sliding window (box filter) to match the 50-ms window used for the Fano factor analysis. Similar results were obtained with a shorter (5-ms) or longer (100-ms) window. (a) Data from one example neuron. V_m for individual trials (black) is plotted on top of the mean (gray). Data are shown when no stimulus was delivered, for a nonpreferred stimulus and for a preferred stimulus. The arrow marks stimulus onset. (b) Similar plot for a second example neuron. (c) The mean and variance of V_m across all 52 neurons and all stimuli. Flanking traces give s.e.m.

Figure 3 Changes in firing-rate variability for ten datasets (one per panel). Insets indicate stimulus type. Data are aligned on stimulus onset (arrow). For the two bottom panels (MT area/direction and MT speed), the dot pattern appeared at time zero (first arrow) and began moving at the second arrow. The mean rate (gray) and the Fano factor (black with flanking s.e.) were computed using a 50-ms sliding window. For OFC, where response amplitudes were small, a 100-ms window was used to gain statistical power. Analysis included all conditions, including nonpreferred. The Fano factor was computed after mean matching (Fig. 4). The resulting stabilized means are shown in black. The mean number of trials per condition was 100 (V1), 24 (V4), 15 (MT plaids), 88 (MT dots), 35 (LIP), 10 (PRR), 31 (PMd), 106 (OFC), 125 (MT direction and area) and 14 (MT speed).



window over the duration of the trial. The Fano factor has been used extensively to characterize neural variability (for example, see refs. 17–19). The Fano factor is influenced both by variability arising from spiking noise and by across-trial variability in the underlying firing rate²⁰. Most prior work assumes that the underlying firing rate is similar across trials and uses the Fano factor to assess the statistics of spiking noise, which are roughly Poisson (Fano factor ≈ 1) for most of cortex. We began with the assumption that spiking noise is roughly Poisson and we used the Fano factor to assess across-trial variability in the underlying rate. We interpreted a Fano factor greater than 1 as being an indication of across-trial firing-rate variability. We interpreted changes in the Fano factor as reflecting changes in across-trial firing-rate variability^{9,20,21}. Although this approach assumes Poisson spiking noise, it is reasonably robust to violations of that assumption (it is sufficient that spiking-noise variance scale linearly with the mean; the slope needn't be unity). To begin, we examined how the Fano factor behaves across a variety of cortical areas.


We computed the mean firing rate and the Fano factor for ten datasets from seven cortical areas of the macaque monkey (Fig. 3): V1, V4, MT, the lateral intra-parietal area (LIP), the parietal reach region (PRR), dorsal premotor cortex (PMd) and orbitofrontal cortex (OFC). Responses were to various visual stimuli or, for OFC, to juice reward. For each area, the Fano factor was averaged across neurons and conditions. This is similar to what was done for the membrane potential analysis and reflects both a desire for statistical power and the expectation that variability may change for both preferred and nonpreferred stimuli (as in Fig. 2a,b).

In every case, stimulus onset drove a decline in firing-rate variability as assessed by the Fano factor (all $P < 0.02$). This is notable, given the diversity of areas, stimuli and behavioral states. Variability declined during responses to simple visual stimuli, during operantly

conditioned responses (PRR and PMd) and during reward-driven responses (OFC). The variability decline was present regardless of whether the monkey was anaesthetized (V1 and two of the four the MT datasets; Fig. 3, bottom), passively viewing (V4) or performing a task (the other six datasets). For two of the MT datasets (Fig. 3, bottom), stimulus onset occurred in two stages: pattern onset and motion onset. Both events drove a decline in variability, although only the more effective moving stimulus drove a sustained decline.

We previously proposed that declining variability in premotor cortex is related to the progress of motor preparation⁹. The changes

Figure 4 Illustration of how the mean-matched Fano factor was computed. Data are from the MT plaids dataset. **(a)** Spike rasters for the 46 trials (one per line) recorded from one MT neuron (127) for one stimulus condition (upwards-moving plaid). Shaded areas show four locations of the sliding window, which moved in 10-ms increments. For each window location, the spike count was computed for each trial. The mean and variance (across trials) of that count then contributed one data point to the subsequent analysis. **(b)** The Fano factor was computed from scatter plots of the spike-count variance versus mean. Each scatter plot corresponds to a window in **a**. Each point plots data for one neuron and condition (red indicates the neuron and condition from **a**). The orange line has unity slope, the expected variance-to-mean relationship for Poisson spiking. Data above the orange line is consistent with the presence of underlying-rate variability. Gray dots show all data. Gray lines are regression fits to all data (constrained to pass through zero, weighted according to the estimated s.e.m. of each variance measurement). Gray distributions are of mean counts. These appear to have different areas because of the vertical log scale. Black points are those preserved by mean matching (Online Methods). Black distributions are thus identical to within bin resolution. Black lines are regression slopes for the mean-matched data. **(c)** The Fano factor versus time. Arrows indicate time points from the panels above. The gray trace (with flanking s.e.) plots the raw Fano factor, the slope of the gray lines from **b**. The black trace plots the mean-matched Fano factor, the slope of the black lines.



in V1 membrane potential variability have been linked to shunting inhibition³ and contrast-invariant orientation tuning⁴. These explanations are no less likely given the current data. However, it is clear that the decline in variability is not limited to those cases but constitutes a cortex-wide effect. This suggests a very general property of cortex, that its state is stabilized by an input.

The mean-matched Fano factor controls for rising rates

The Fano factor declines as mean firing rates rise. This was expected; in the intracellular data, it was similarly true that V_m variability dropped as the mean V_m rose. It is nevertheless important to rule out any possibility that the declining Fano factor is trivially related to rising rates. At higher rates, refractory periods will tend to regularize spiking (a violation of the Poisson assumption), which could reduce the Fano factor simply because the variability of spiking-process noise is reduced. A less obvious concern is that across-trial firing-rate variability could be constant but becomes normalized by a higher mean after stimulus onset (the Fano factor is the spike-count variance divided by the mean).


To address these issues, we computed the Fano factor traces (Fig. 3) using a method that we refer to as mean matching. Interpretation would be aided if we could record from a population of neurons that did not change its overall distribution of mean firing rates. If some neurons responded to the stimulus with an increased mean rate and others with a decreased mean rate, then the overall distribution of mean rates (across neurons and conditions) could stay conveniently constant. For many areas (for example, PRR), this is not

far from what was observed. The overall mean rate always increased poststimulus, but this was often driven by a small percentage of neurons that strongly preferred the stimulus. Most neurons exhibited more modest changes in mean rate, with declines being almost as common as increases. For other areas (especially anesthetized V1), it was the case that the vast majority of neurons increased their mean rate following stimulus onset. Nevertheless, for all datasets, there was considerable overlap in the distribution of mean rates before and after stimulus onset. Mean matching exploits this overlap.

To apply mean matching, we computed the mean and variance of the spike count in a sliding window for each neuron and condition (Fig. 4a). At each time, we plotted the variances versus their respective means, with one data point for every neuron and condition (Fig. 4b). The raw Fano factor was simply the slope of the regression relating the variance to the mean. The concern being addressed is that the raw Fano factor might be lower following stimulus onset simply because rates are higher. Inspection revealed this to be unlikely; even for the range of mean counts that is shared across times, there were noticeably fewer high-variance points for the later times (compare -100 and +100 ms). Mean matching formalized this observation. We computed distributions of mean counts and took the greatest common distribution. At each time, individual points were randomly excluded until the actual distribution matched the common distribution. The mean-matched Fano factor was based on the remaining points.

All of the Fano factors in Figure 3 employed mean matching. Very similar results were obtained for the raw Fano factor (as in Fig. 4c). Thus, the decline is not somehow produced by mean matching. Mean

Figure 5 Changes in the Fano factor after restricting the analysis to combinations of neuron and condition with little change in mean rate (for example, nonpreferred conditions). **(a–d)** The raw Fano factor for four datasets, computed based on nonresponsive conditions. Of the original neuron conditions (the response of one neuron to one condition), this analysis preserved 28% (MT, **a**), 49% (PRR, **b**), 27% (PMd, **c**) and 41% (MT-speed, **d**). A 100-ms window (rather than 50 ms) was employed to regain lost statistical power. The trace at the top of each panel shows the mean rate, averaged across all included neurons and conditions. The trace with flanking s.e. shows the Fano factor, computed with no further mean matching. Arrows indicate stimulus onset. For the MT speed dataset (**d**) the stimulus appeared at the very start of the record (first arrow) and began moving 256 ms later (second arrow).



matching simply controls for a potential artifact. Also note that most of the analyzed neurons still showed stimulus-driven changes in mean rate (both increases and decreases), even following mean matching. Only the overall distribution of firing rates was held constant. Before application to real data, mean matching was applied to many simulated datasets (a few of which are shown in **Supplementary Fig. 2**). The mean-matched Fano factor successfully avoids artifacts related to changing mean rates over a wide range of circumstances, including low trial counts, non-Poisson spiking, low mean rates and large changes in mean rate.

Further controls for the Fano factor

We employed a complementary method to further control for potential changes in spiking-noise statistics. We computed both the Fano factor (for all neurons and conditions) and the square of the coefficient of variation of the interspike intervals using a previously described method²⁰. Although the Fano factor declined following stimulus onset, the square of the coefficient of variation changed little (**Supplementary Fig. 3**). This is not consistent with a large change in spiking-process statistics (which should produce a large decline in both the Fano factor and the square of the coefficient of variation) but is consistent with a decline in across-trial rate variability, a similar conclusion to that drawn previously²⁰. One can also control for rising mean rates (and resulting refractoriness) by focusing on conditions and neurons that individually exhibit little change in mean rate (for example, null directions and other non-responsive conditions). This necessitates that there are a sufficient number of such cases to allow further analysis. Of the datasets in **Figure 3**, four (MT plaids, PRR, PMd and MT speed) could be analyzed in this manner. In each case (**Fig. 5**), the Fano factor decline was as clear for the nonresponsive conditions ($P < 10^{-6}$, $P < 10^{-6}$, $P < 0.002$ and $P < 10^{-4}$) as it had been for the original analyses.

This finding also addresses a second potential concern: low-rate neurons (which are preferentially preserved by mean matching following stimulus onset) might tend to have low intrinsic Fano factors, resulting in an artifactual Fano factor decline. However, such an artifact cannot affect the present analysis (**Fig. 5**); exactly the same neurons and conditions contributed at all of the times. Furthermore, the raw Fano factor always showed the same decline as the mean-matched Fano factor. In summary, the decline in the Fano factor

cannot be a simple consequence of refractoriness, changing spiking-noise statistics or normalization. Instead, it appears to reflect a widespread decline in the variance of the underlying firing rate.

Variability declines even for nonresponsive conditions

Analyses of nonresponsive conditions (**Figs. 2 and 5**) acts as a control but also argues for an additional point. The firing rate, averaged across trials, may change little, but neurons are not truly nonresponsive. It appears that underlying rates do change from a variable range of rates before stimulus onset to a more consistent rate (with a similar mean) following stimulus onset. It may be only the mean rate that changes little (**Fig. 1c**).

The critical factor is not whether a neuron appears to be responsive from its mean, but whether the population as a whole is responsive. This is illustrated by our data from anesthetized MT (**Fig. 3**, bottom). The onset of a static dot pattern evoked a weak and transient increase in mean rate at the population level and a transient (though large) decline in the Fano factor. In contrast, dot motion onset evoked a sustained increase in the mean population rate and a sustained Fano factor decline. This Fano factor decline was present and sustained even for those neurons where the mean rate changed little (**Fig. 5d**). Thus, the Fano factor decline depends on the stimulus effectively driving the population as a whole, but not on individual neuron mean responsiveness. Consistent with this, the Fano factor rose again fairly rapidly when the population ceased to respond following stimulus offset (**Supplementary Fig. 4**).

A decline in correlated variability

Does each neuron undergo its own decline in variability or is the above effect related to variability that is shared? To assess this, we exploited the covariance of simultaneous recordings, obtained via 96-electrode array recordings from V1 and PMd. The measured covariance matrix (assessed across trials) reflects both shared firing-rate variability and 'private' spiking-process noise. It is thus necessary to decompose the measured covariance matrix into shared (network level) and private components (**Fig. 6a**). A number of applicable methods exist, all of which exploit the assumption that shared variability should be relatively low dimensional (that is, that neurons are

tuned to a modest number of shared factors). Intuitively, if one can average across similarly tuned neurons to remove spiking noise, the underlying rate (and its variance) becomes more accessible. More precisely, if neurons are tuned for eight shared factors, the covariance matrix describing the underlying rate will be of rank 8. The measured covariance matrix should then be decomposed into a rank-8 'network' covariance matrix and a diagonal matrix capturing private spiking-process noise. Spiking-process variability is not necessarily physiologically cell intrinsic^{14,16,17} but is weakly correlated between most neurons and thus private for our purposes, especially given the long analysis windows being used. Of the applicable methods²², factor analysis²³ has the most appropriate noise model; different neurons may have different levels of private noise. This is essential, as Poisson spiking noise scales with the mean rate (how factor analysis separates network variability from private spiking noise is shown in **Supplementary Fig. 5**).

We applied factor analysis to four datasets: two V1 datasets from different monkeys and two PMd datasets collected from one monkey on separate days (**Fig. 6**; see Online Methods and **Supplementary Fig. 6**

for a description of datasets). After isolating the network variances (**Fig. 6a**), we computed the average network variance (across neurons and conditions) before and after stimulus onset. The average variance was computed following a mean-matching procedure, just as was performed for the Fano factor. Mean matching aids interpretation by allowing direct comparison of the variance present for a given mean rate. It also insures that any contamination of network variability by private noise (inevitable for finite trial counts) does not increase with rising rates.

A decline in network variability was observed in all of the datasets (**Fig. 6b,c,e,f**). For V1, network variability was 66% and 40% lower following stimulus onset. For PMd, network variability was 53% and 45% lower. This decline was significantly greater for network variability than for private noise, in both absolute and relative terms (**Fig. 6d,g**). Thus, the Fano factor and factor analyses agree that stimulus onset drives a decline in firing-rate variability that is not simply the result of a change in the regularity of spiking itself.

The decline in rate covariance is consistent with a previous study²⁴ (from which one of the V1 datasets is drawn), in which correlations

Figure 6 Application of factor analysis to data from V1 and PMd. **(a)** Factor analysis was applied to covariance matrices (number of neurons \times number of neurons) of spike counts, taken in an analysis window that either ended at stimulus onset (prestimulus) or began just after stimulus onset (stimulus). The measured covariance matrix was approximated as the sum of a network covariance matrix and a diagonal matrix of private noise. To produce the plots in **b–g**, we averaged network variances across the subset of neuron and condition combinations (48% and 30% for V1, 74% and 79% for PMd) whose distribution of mean rates was matched before and after stimulus onset (similar to **Fig. 4**). **(b)** Estimated variances for one V1 dataset. Network variability declined more than private variability in both absolute ($P < 10^{-7}$) and relative (percent of initial value, $P < 10^{-7}$) terms (paired t tests across conditions). **(c)** Similar plot for a V1 dataset from a second monkey ($P < 0.002$, absolute; $P < 0.002$, relative). **(d)** Summary comparison for V1. Changes in variability (stimulus–prestimulus) were expressed in percentage terms. Data to the left of zero indicate that network variability underwent the larger decline. The distribution includes all conditions and both datasets. The mean and s.e. are given by the black symbol at top ($P < 10^{-7}$ compared with zero, paired t test). Gray symbols give individual means for each dataset. **(e)** Data are presented as in **b** and **c** but for one PMd dataset (G20040123). Network variability declined more in absolute ($P < 0.005$) and relative ($P < 0.001$) terms. **(f)** Similar plot for a second PMd dataset (G20040122; $P < 0.05$, absolute; $P < 0.02$, relative). **(g)** Summary comparison for PMd (distribution mean < 0 , $P < 10^{-4}$). **(h)** Relationship between mean firing rate and network level (shared firing rate) variance. Data (same dataset as **b**) were binned by mean rate and the average network variance (\pm s.e.) was computed for each bin. This was done both before stimulus (gray) and after stimulus onset (black). The average was taken across neurons and conditions (each datum being averaged was, for one condition, one element of the blue diagonal in **a**). Distributions of mean rates are shown at bottom. The analysis in **b** was based on the overlapping (mean matched) portion of these distributions. **(i)** Similar plot for PMd (same dataset as **e**). See Online Methods and **Supplementary Figure 6** for a description of datasets.





Figure 7 Individual-trial neural trajectories computed using GPFA.

(a) Projections of PMd activity into a two-dimensional state space. Each black point represents the location of neural activity on one trial. Gray traces show trajectories from 200 ms before target onset until the indicated time. The stimulus was a reach target (135°, 60 mm distant), with no reach allowed until a subsequent go cue. 15 (of 47) randomly selected trials are shown. The dataset is the same as in **Figure 6e**. (b) Trajectories were plotted until movement onset. Blue dots indicate 100 ms before stimulus (reach target) onset. No reach was allowed until after the go cue (green dots), 400–900 ms later. Activity between the blue and green dots thus relates to movement planning. Movement onset (black dots) was ~300 ms after the go cue. For display, 18 randomly selected trials are plotted, plus one hand-selected trial (red, trialID 211). Covariance ellipses were computed across all 47 trials. This is a two-dimensional projection of a ten-dimensional latent space. In the full space, the black ellipse is far from the edge of the blue ellipse. This projection was chosen to accurately preserve the relative sizes (on a per-dimension basis) of the true ten-dimensional volumes of the ellipsoids. Data are from the G20040123 dataset. (c) Data are presented as in **b**, with the same target location, but for data from another day's dataset (G20040122; red trial, trialID 793).



were lower overall after stimulus onset. However, our result does not simply follow from that result. The mean pair-wise correlation can decline even as overall covariance grows (for example, if negative correlations become more common or if private spiking variability climbs).

In three of the four cases, factor analysis also indicated a small reduction in private noise. Limited data make it difficult to determine whether this indicates an actual change in spiking-process statistics or if it occurs because some network variability is incorrectly assigned to private noise. The latter can occur if the true dimensionality of the data is higher than the dimensionality assumed by the analysis (**Supplementary Fig. 7**). This possibility is of minor concern, as it would imply that we are underestimating the network-variance decline. The converse (mis-assignment of a private change to a network change) is potentially concerning, but unlikely; for trial-shuffled data, factor analysis correctly assigns the variance change to private noise with only a small leak into network variance (**Supplementary Fig. 7**). Furthermore, the decline in network variability was the larger effect, both in absolute (**Fig. 6b,c,e,f**) and percentage (**Fig. 6d,g**) terms. In summary, there may or may not be modest changes in private spiking-process variability following stimulus onset. In either case, the measured decline in variability is principally a decline in network-level variability that is shared among neurons.

As an aside, the shared variability being assessed is indeed variability in the firing rate across trials and not variability in precise spike patterns. Recomputing the Fano factor and factor analysis after 'jittering' each spike in a 20-ms window produced essentially identical results. This was unsurprising, given the large (400 ms) analysis window.

A qualification regarding the decline in variability

For both the Fano factor and for factor analysis, we compared identical distributions of mean rates before and after stimulus onset. We found that for a given mean rate there was less variability around that mean following stimulus onset (for example, the variance around a 10 spike per s mean might shrink from 15 to 8 spikes² per s²). This will often imply that overall variability declines, but this may not occur when rates are initially very low. In the limit, when initial firing rates are zero, variability is zero and will necessarily rise when rates rise. For area PMd, overall firing rates changed modestly following

stimulus onset and overall variability did indeed decline. In anaesthetized V1, however, most neurons had baseline rates near zero and therefore had near-zero variability. Overall variability is therefore essentially guaranteed to rise following stimulus onset. This can be seen by plotting network-level rate variability versus mean rate (**Fig. 6h,i**). In this analysis, spiking variability was approximately factored out, leaving primarily underlying-rate variability. The effect of stimulus onset was essentially the same for both PMd and V1. For any given mean rate, there was less variability around that rate after stimulus onset. For V1, however, mean rates were low before stimulus onset, necessitating low rate variance. Overall variability therefore rose following stimulus onset (average variance rose from 77 ± 5 s.e.m. to 190 ± 8 spikes² per s²) even though it was lower for any given mean rate. For PMd, the overall change in mean rates was modest and variance was lower after stimulus onset both for a given rate and in overall terms (average variance dropped from 28 ± 2 to 16 ± 1 spikes² per s²).


Thus, the central effect is a decline in the variability present around a given mean rate. This may or may not translate into an overall variance decline, depending on whether overall mean rates rise modestly or sharply. This qualification is restricted to the analysis of firing rate, whose variability must be zero when the mean is zero. In contrast, the membrane voltage showed considerable variability even when firing rates were near zero (**Fig. 2**).

Single trial analysis of variability

The neural response is incompletely characterized by its mean. A fuller characterization is provided by also assessing firing-rate variance. Ideally, however, one would directly observe the evolution of firing rates on individual trials, much as we directly observed the membrane potential on individual trials (**Fig. 2**). Such visualization is likely important, not only for the initially high variance but also for the variance that survives stimulus onset. The presence of remaining firing-rate variability after stimulus onset is underscored by its correlation with pre-stimulus variability^{2,6,8} (**Supplementary Note 1**). Factor analysis (**Fig. 6h,i**) also indicated that considerable firing-rate variability remains following stimulus onset. For PMd, such variability is behaviorally meaningful: it correlates with subsequent reaction time⁹ and reach speed²⁵.

Estimating the evolution of single-trial firing rates is difficult or impossible from single-neuron recordings, but becomes plausible

Figure 8 Projections of V1 activity into a two-dimensional space using GPFA. Blue, black and red traces show activity before, during and after stimulus presentation (a drifting 45° grating). Data are from the dataset used in **Figure 6c**. **(a)** The mean trajectory and three trials picked by hand. The gray spot shows the average location of prestimulus activity. In a few cases (for example, upper left portion of the rightmost panel), traces were moved very slightly apart to make it clear that they traveled in parallel rather than crossed. **(b)** Trajectories after data were shuffled to remove correlated variability. 25 randomly selected trials are plotted (lighter traces) along with the mean (saturated traces). **(c)** Data are presented as in **b** but for the original unshuffled data.



for large-scale simultaneous recordings. To do so, we employed an extension of factor analysis termed Gaussian-process factor analysis (GPFA)²⁶. GPFA projects the responses of many neurons into a low-dimensional space where each trial's neural trajectory can be traced through time. Essentially, each axis captures the smoothed and averaged response of similarly tuned neurons. More precisely, each neuron's underlying rate is a linear combination of the responses represented on each axis (much as for principal component analysis). A full characterization of single-trial variability across all conditions was beyond the scope of our study. Still, our analyses serve to illustrate the power and, in some cases, the limitations of viewing neural responses on a single-trial basis.

The collapse in variance in PMd was readily visualized via GPFA (**Fig. 7a**). The neural state was initially variable across trials, even though the monkey's external state, including fixation, was tightly controlled. Following reach-target onset, the neural state became less variable and changed its center of mass (that is, mean rates changed). Across both datasets and all 14 target locations, variability was always lower by the go cue than it had been pre-stimulus (for the 2 datasets, variance was an average of 27% and 37% smaller per dimension, relative to 100 ms pre-stimulus, $P < 10^{-6}$ and $P < 10^{-8}$; t tests across 14 conditions). Variability was lower still by movement onset (36% and 46% smaller than pre-stimulus variability; $P < 10^{-10}$, $P < 10^{-11}$). These features are particularly clear in video format (**Supplementary Videos 1–4**).

Despite these results, the principal goal of GPFA is not to further quantify variability and its decline (that is better accomplished by the Fano factor and by factor analysis). Rather, we would like to know how individual trials deviate from their mean. In particular, although most neural trajectories hewed close to their mean after target onset, there were exceptions. For example, one trial followed the usual trajectory up until the go cue (presumably indicating successful target detection and initial planning; **Fig. 7b**). Following the go cue, the trajectory wandered before falling back on track. On this trial, the monkey had an unusually long reaction time (629 ms, all others ranged from 240–375 ms). Other trials, for other target locations and days, contained similar outliers associated with long reaction times (**Fig. 7c**, **Supplementary Fig. 8** and **Supplementary Videos 1–4**). Thus, across-trial averaging obscures not only the variance decline but also other features, including events that may only have happened once.

We also applied GPFA to data from V1. We examined neural trajectories for individual trials (data for one condition, employing a 45° drifting grating; **Fig. 8**). The initial overshoot corresponds to the onset transient and the looping trajectory was driven by the periodic stimulus. Across-trial variability was, if anything, more pronounced than in PMd. Individual trials showed considerable variability in the stimulus-driven trajectory and in the trajectory both before stimulus onset and after its offset. Of course, given a limited number of neurons, GPFA cannot provide noise-free estimates of single-trial firing rates. However, much of the observed variability was real; shuffling the data to break neuron-neuron correlations greatly reduced

variability relative to the original data (**Fig. 8b,c**). This was similarly true for every stimulus condition for all of the datasets tested. For the two V1 datasets, the original data had, on average, 81% and 89% more variance ($P < 10^{-10}$ for both, t test across conditions) than the shuffled data. The same was true for PMd (42% and 43% more variance, $P < 10^{-15}$ for both). Thus, the variable individual trial trajectories do indeed reflect trial-to-trial departures from the mean behavior. In particular, for V1 there was a notable degree of variability in the initial state (**Fig. 8c**), especially given the low average initial rates of most neurons. Because mean matching cannot be applied to GPFA, total variability certainly did not decline following stimulus onset. Indeed it was higher, consistent with the results of factor analysis (**Fig. 6h**). In summary, GPFA is in some ways less flexible than the Fano factor or factor analysis (mean matching cannot be applied and the noise model is fixed across time). Still, the ability to reconstruct single-trial responses and observe events that may have occurred only once lends it a unique utility.

DISCUSSION

Stimulus onset drives a widespread decline in the variability of the neural state. Consistent with prior reports, this effect was directly observable at the level of the membrane potential^{3,4,27}. At the level of firing rates, the effect was not initially obvious. Much of the measured variability at the level of a single neuron was the result of quasi-Poisson spiking statistics, whose variance rises with mean rate¹⁷. Nevertheless, appropriately controlled metrics revealed that firing-rate variability declined sharply following stimulus onset. Specifically, for a given mean rate, there was much less variability around the rate after stimulus onset. The ability to assess this effect in extracellular recordings allowed us to examine datasets from many cortical areas, employing many stimuli, in both anesthetized and behaving monkeys. Stimulus onset might have been expected to affect neural variability differently depending on the area, stimulus, task and behavioral state. However, we consistently observed a decline in neural variability across all 14 datasets, from occipital to frontal cortex. This finding argues that a stimulus-driven decline in variability is a widespread feature of the cortical response.

The Fano factor and factor analysis both indicate that the central effect is a decline in the across-trial variability of the underlying

firing rate, rather than a change in the spiking-statistics applied to that rate. This decline in firing-rate variability is probably a network property, for two reasons. First, the decline occurred even when the mean firing rate of a given neuron changed little (for example, for non-preferred stimuli), provided that the network as a whole responded to the stimulus. Second, the variability in question was shared among neurons in an area, as revealed by factor analysis. This finding, and the similar decline across areas, suggests that the relevant network could span much of the cortex. Alternately, it may be that different areas exhibit the same effect simply because it is a common property of cortex. Distinguishing these possibilities will require large-scale simultaneous recordings from multiple areas.

Many prior studies have examined correlations between firing-rate variability and behavior (for example, ref. 28). A few have related across-trial firing-rate variability with task difficulty²⁹, attention^{21,30} or motor learning^{31,32}. Our study is, to the best of our knowledge, the first to fully document a simpler effect: the change in variability driven by the stimulus itself. However, related effects can certainly be found in prior studies. This is true not only for the membrane potential^{3,4} but also for firing-rate variability. We previously reported (using methodology related to the Fano factor) a variability decline in premotor cortex⁹. Variability also undergoes changes in motor cortex^{10,20,33}. In retina, LGN and V1, the Fano factor was lowest when rates were highest³⁴. In a recent study of V4 (ref. 21), the Fano factor dropped after the stimulus entered the receptive field. There is also a small Fano factor drop in previously published MT data³⁵. Variability declined throughout the trial for multiple areas during a probabilistic choice task³⁶. Both cerebellar and FEF variability reach minima before movement^{37,38}. A stimulus-driven decline in correlated activity^{5,24} was found for V1, consistent with a decline in (correlated) rate variability (also see ref. 39 for a similar effect in the LFP). Finally, there are reports⁴⁰ of lower variability in driven versus spontaneous thalamic activity. In some of these studies, variability changes were of passing interest and controls for changing spiking-process statistics were not always applied (although see refs. 9,20). However, our results argue that most of these effects probably reflect a decline in network-level firing-rate variance.

Mechanistically, the variance decline implies that cortical circuits become more stable when driven. Many network types can be stabilized by an input. Natural candidates are recurrent networks with attractor dynamics⁴¹. More broadly, a stimulus-driven decline in variability may be a general feature of large recurrent networks^{11,12}. The widespread effects that we observed may simply reflect the widespread presence of recurrent circuitry. These network-level explanations are consistent with the shared nature of the firing-rate variability and with our previous finding that a drop in PMd variability relates to reaction time⁹. However, other data indicate a bottom-up source; in particular, shunting inhibition appears to be the source of the decline in V1 membrane-potential variability³. One hopes that these two levels of explanation will be reconciled, although it is not yet clear how to do so.

The functional interpretation of converging rates likely varies by area. For PMd, the decline may relate to motor preparation^{9,10}. In area V4, it may relate to attention²¹. Often it is tempting to relate the variability decline to cognitive-level phenomena; stimulus onset is a natural candidate to focus not only attention and planning, but other forms of mental processing as well. On the other hand, the decline was observed even under anesthesia (all V1 datasets and two of the MT datasets), consistent with lower-level explanations³ (see **Supplementary Note 2** for further discussion of the possible origins, local versus cognitive, of the initial variability). The anesthetized data also rule out effects related to microsaccades or other unobserved changes in behavior. Finally,

effects in the anesthetized data appear unrelated to long timescale drift of anesthesia; the decline was undiminished when computed more locally in time (**Supplementary Note 3**).

Despite being widespread, the changes in variability are unlikely to constitute a coding channel. It is unlikely that the brain could measure or interpret variability that is present only across trials. It is the experimenter who, when interpreting the recorded responses, needs to know when variability is substantial and how it is changing. Fundamentally we need to know when the mean is representative. Notably, different classes of dynamics (single-attractor, multi-attractor and integrator) can have different variability 'signatures' (A.K. Churchland, R. Kiani & M.N. Shadlen, *Soc. Neurosci. Abstr.* **548.29**, 2006). Variability measurements may thus allow hypotheses regarding circuit-level dynamics to be tested at the level of the single neuron. More generally, although the mean and the variance are indispensable metrics, one also wishes to observe the individual-trial responses themselves. GPFA and related methods can approximately reconstruct individual-trial neural trajectories from simultaneous extracellular recordings⁴² and thus have the potential to uncover telling features of the neural response that are normally lost to averaging.

METHODS

Methods and any associated references are available in the online version of the paper at <http://www.nature.com/natureneuroscience/>.

Note: Supplementary information is available on the Nature Neuroscience website.

ACKNOWLEDGMENTS

This work was supported by a Helen Hay Whitney postdoctoral fellowship (M.M.C.), Burroughs Wellcome Fund Career Awards in the Biomedical Sciences (M.M.C. and K.V.S.), Gatsby Charitable Foundation (M.S. and B.M.Y.), US National Institutes of Health (NIH), National Institute of Neurological Disorders and Stroke Collaborative Research in Computational Neuroscience grant R01-NS054283 (K.V.S. and M.S.), the Michael Flynn Stanford Graduate Fellowship (J.P.C.), the Howard Hughes Medical Institute and NIH grant EY05603 (W.T.N., L.P.S., M.R.C. and G.S.C.), a Howard Hughes Medical Institute predoctoral fellowship (M.R.C.), NIH grant EY014924 (T.M. and K.M.A.), Sloan Foundation (T.M. and K.V.S.), Pew Charitable Trust (T.M.), NIH EY015958 and EY018894 (M.A.S.), NIH EY02017 and EY04440 (J.A.M.), NIH EY016774 (A.K.), NIH 1 EY13138-01 (D.C.B., A.M.C., P.H. and B.B.S.), NIH EY019288 (N.J.P.), the Pew Charitable Trust (N.J.P.), EY04726 (D.F.), US National Defense Science and Engineering Graduate Fellowships (B.M.Y. and G.S.), National Science Foundation Graduate Research Fellowships (B.M.Y. and G.S.), the Christopher and Dana Reeve Foundation (K.V.S. and S.I.R.), and the awards from Stanford Center for Integrated Systems, National Science Foundation Center for Neomorphic Systems Engineering at Caltech, Office of Naval Research, NIH Director's Pioneer Award 1DP1OD006409 and Whitaker Foundation (K.V.S.).

AUTHOR CONTRIBUTIONS

M.M.C. wrote the manuscript, performed the Fano factor and factor analyses and created the figures. GPFA was developed by B.M.Y., J.P.C., M.S. and K.V.S. This application of factor analysis was devised by M.M.C. and B.M.Y. The mean-matched Fano factor was developed by M.M.C. and K.V.S. The conception for the study arose from conversations between M.M.C., K.V.S., B.M.Y., D.C.B., M.R.C., W.T.N. and J.A.M. V1 data (extracellular) were collected in the laboratory of J.A.M. by M.A.S. and A.K. and in the laboratory of A.K. V4 data were collected in the laboratory of T.M. by K.M.A. MT (plaid) data were collected in the laboratory of D.C.B. by A.M.C., P.H. and B.B.S. MT (dots) data were collected in the laboratory of W.T.N. by M.R.C. LIP and OFC data were collected in the laboratory of W.T.N. by L.P.S. using an experimental design developed by L.P.S. and G.S.C. PRR data were collected in the laboratory of L.H.S. by S.W.C. PMd data were collected in the laboratory of K.V.S. by B.M.Y., S.I.R., G.S. and M.M.C. MT (direction/area and speed) data were collected by N.J.P. and M.M.C. in the laboratory of S.G.L. Intracellularly recorded V1 data were collected by N.J.P. and I.M.F. in the laboratory of D.F. All authors contributed to manuscript revisions and editing, particularly J.A.M., W.T.N., L.P.S., D.F., J.P.C., B.M.Y. and K.V.S.

COMPETING INTERESTS STATEMENT

The authors declare no competing financial interests.

Published online at <http://www.nature.com/natureneuroscience/>.

Reprints and permissions information is available online at <http://www.nature.com/reprintsandpermissions/>.

1. Briggman, K.L., Abarbanel, H.D. & Kristan, W.B. Jr. Optical imaging of neuronal populations during decision-making. *Science* **307**, 896–901 (2005).
2. Arieli, A., Sterkin, A., Grinvald, A. & Aertsen, A. Dynamics of ongoing activity: explanation of the large variability in evoked cortical responses. *Science* **273**, 1868–1871 (1996).
3. Monier, C., Chavane, F., Baudot, P., Graham, L.J. & Fregnac, Y. Orientation and direction selectivity of synaptic inputs in visual cortical neurons: a diversity of combinations produces spike tuning. *Neuron* **37**, 663–680 (2003).
4. Finn, I.M., Priebe, N.J. & Ferster, D. The emergence of contrast-invariant orientation tuning in simple cells of cat visual cortex. *Neuron* **54**, 137–152 (2007).
5. Kohn, A., Zandvakili, A. & Smith, M.A. Correlations and brain states: from electrophysiology to functional imaging. *Curr. Opin. Neurobiol.* **19**, 434–438 (2009).
6. Azouz, R. & Gray, C.M. Cellular mechanisms contributing to response variability of cortical neurons *in vivo*. *J. Neurosci.* **19**, 2209–2223 (1999).
7. Fiser, J., Chiu, C. & Weliky, M. Small modulation of ongoing cortical dynamics by sensory input during natural vision. *Nature* **431**, 573–578 (2004).
8. Kisley, M.A. & Gerstein, G.L. Trial-to-trial variability and state-dependent modulation of auditory-evoked responses in cortex. *J. Neurosci.* **19**, 10451–10460 (1999).
9. Churchland, M.M., Yu, B.M., Ryu, S.I., Santhanam, G. & Shenoy, K.V. Neural variability in premotor cortex provides a signature of motor preparation. *J. Neurosci.* **26**, 3697–3712 (2006).
10. Rickert, J., Riehle, A., Aertsen, A., Rotter, S. & Nawrot, M.P. Dynamic encoding of movement direction in motor cortical neurons. *J. Neurosci.* **29**, 13870–13882 (2009).
11. Sussillo, D. & Abbott, L.F. Generating coherent patterns of activity from chaotic neural networks. *Neuron* **63**, 544–557 (2009).
12. Abbott, L.F., Rajan, K. & Sompolinsky, H. Interactions between intrinsic and stimulus-dependent activity in recurrent neural networks. in *Neuronal Variability and its Functional Significance* (eds. Ding, M. & Glanzman, D.) (in the press).
13. Poulet, J.F. & Petersen, C.C. Internal brain state regulates membrane potential synchrony in barrel cortex of behaving mice. *Nature* **454**, 881–885 (2008).
14. Shadlen, M.N. & Newsome, W.T. The variable discharge of cortical neurons: implications for connectivity, computation, and information coding. *J. Neurosci.* **18**, 3870–3896 (1998).
15. van Vreeswijk, C. & Sompolinsky, H. Chaos in neuronal networks with balanced excitatory and inhibitory activity. *Science* **274**, 1724–1726 (1996).
16. Mainen, Z.F. & Sejnowski, T.J. Reliability of spike timing in neocortical neurons. *Science* **268**, 1503–1506 (1995).
17. Carandini, M. Amplification of trial-to-trial response variability by neurons in visual cortex. *PLoS Biol.* **2**, e264 (2004).
18. Tolhurst, D.J., Movshon, J.A. & Dean, A.F. The statistical reliability of signals in single neurons in cat and monkey visual cortex. *Vision Res.* **23**, 775–785 (1983).
19. Gur, M., Beylin, A. & Snodderly, D.M. Response variability of neurons in primary visual cortex (V1) of alert monkeys. *J. Neurosci.* **17**, 2914–2920 (1997).
20. Nawrot, M.P. *et al.* Measurement of variability dynamics in cortical spike trains. *J. Neurosci. Methods* **169**, 374–390 (2008).
21. Mitchell, J.F., Sundberg, K.A. & Reynolds, J.H. Differential attention-dependent response modulation across cell classes in macaque visual area V4. *Neuron* **55**, 131–141 (2007).
22. Roweis, S. & Ghahramani, Z. A unifying review of linear gaussian models. *Neural Comput.* **11**, 305–345 (1999).
23. Everitt, B.S. *An Introduction to Latent Variable Models* (Chapman & Hall, London, 1984).
24. Smith, M.A. & Kohn, A. Spatial and temporal scales of neuronal correlation in primary visual cortex. *J. Neurosci.* **28**, 12591–12603 (2008).
25. Churchland, M.M., Afshar, A. & Shenoy, K.V. A central source of movement variability. *Neuron* **52**, 1085–1096 (2006).
26. Yu, B.M. *et al.* Gaussian-process factor analysis for low-dimensional single-trial analysis of neural population activity. *J. Neurophysiol.* **102**, 614–635 (2009).
27. Monier, C., Fournier, J. & Fregnac, Y. *In vitro* and *in vivo* measures of evoked excitatory and inhibitory conductance dynamics in sensory cortices. *J. Neurosci. Methods* **169**, 323–365 (2008).
28. Britten, K.H., Newsome, W.T., Shadlen, M.N., Celebri, S. & Movshon, J.A. A relationship between behavioral choice and the visual responses of neurons in macaque MT. *Vis. Neurosci.* **13**, 87–100 (1996).
29. Horwitz, G.D. & Newsome, W.T. Target selection for saccadic eye movements: prelude activity in the superior colliculus during a direction-discrimination task. *J. Neurophysiol.* **86**, 2543–2558 (2001).
30. Cohen, M.R. & Maunsell, J.H. Attention improves performance primarily by reducing interneuronal correlations. *Nat. Neurosci.* **12**, 1594–1600 (2009).
31. Mandelblat-Cerf, Y., Paz, R. & Vaadia, E. Trial-to-trial variability of single cells in motor cortices is dynamically modified during visuomotor adaptation. *J. Neurosci.* **29**, 15053–15062 (2009).
32. Kao, M.H., Doupe, A.J. & Brainard, M.S. Contributions of an avian basal ganglia-forebrain circuit to real-time modulation of song. *Nature* **433**, 638–643 (2005).
33. Oram, M.W., Hatsopoulos, N.G., Richmond, B.J. & Donoghue, J.P. Excess synchrony in motor cortical neurons provides redundant direction information with that from coarse temporal measures. *J. Neurophysiol.* **86**, 1700–1716 (2001).
34. Kara, P., Reinagel, P. & Reid, R.C. Low response variability in simultaneously recorded retinal, thalamic, and cortical neurons. *Neuron* **27**, 635–646 (2000).
35. Osborne, L.C., Bialek, W. & Lisberger, S.G. Time course of information about motion direction in visual area MT of macaque monkeys. *J. Neurosci.* **24**, 3210–3222 (2004).
36. Lee, D. & Seo, H. Neural and behavioral variability related to stochastic choices during a mixed-strategy game. in *Neuronal Variability and its Functional Significance* (eds. Ding, M. & Glanzman, D.) (in the press).
37. Fortier, P.A., Smith, A.M. & Kalaska, J.F. Comparison of cerebellar and motor cortex activity during reaching: directional tuning and response variability. *J. Neurophysiol.* **69**, 1136–1149 (1993).
38. Cohen, J.Y. *et al.* Difficulty of visual search modulates neuronal interactions and response variability in the frontal eye field. *J. Neurophysiol.* **98**, 2580–2587 (2007).
39. Nauhaus, I., Busse, L., Carandini, M. & Ringach, D.L. Stimulus contrast modulates functional connectivity in visual cortex. *Nat. Neurosci.* **12**, 70–76 (2009).
40. Werner, G. & Mountcastle, V.B. The variability of central neural activity in a sensory system and its implications for the central reflection of sensory events. *J. Neurophysiol.* **26**, 958–977 (1963).
41. Wang, X.J. Decision making in recurrent neuronal circuits. *Neuron* **60**, 215–234 (2008).
42. Churchland, M.M., Yu, B.M., Sahani, M. & Shenoy, K.V. Techniques for extracting single-trial activity patterns from large-scale neural recordings. *Curr. Opin. Neurobiol.* **17**, 609–618 (2007).

ONLINE METHODS

Data sets. All methods were approved in advance by the relevant institutional animal care and use committees. Data were collected from seven cortical areas of the macaque monkey and one cortical area of the cat by members of nine laboratories. For most datasets, methods have been described previously^{4,9,24,43–48}. Briefly, for the intracellularly recorded V1 datasets, stimuli were 100% contrast sine-wave gratings presented at different orientations. For the extracellularly recorded V1 dataset, stimuli were 100% contrast sine-wave gratings (6.25 Hz, 1.3 cycles per degree) drifting in 1 of 12 directions. For V4, stimuli were one or two oriented bars situated in the neuron's receptive field. For some conditions, similar bars appeared in the opposite hemifield. For the first MT dataset, stimuli were square-wave gratings superimposed to produce a plaid. These patterns contained a small amount of variable-contrast texture, which was identical for all trials of a given type in a recording session (making it sensible to compute the response variance across those trials). For the second MT dataset, stimuli were 0% coherence random dots. These had exactly the same dot pattern for repeated trials in a session. For LIP, stimuli were two colored saccade targets, between which the monkey chose. Analysis was further segregated by the eventual choice and the estimated value of that choice to the monkey. For example, one of the conditions corresponded to rightwards saccades to a red target of high reward probability. For PRR, stimuli were reach and saccade targets (modality instructed by color) both inside and outside of the neuron's receptive field. For PMd, stimuli were reach targets at one of two distances and seven directions. For OFC, the task was similar to that used for LIP, but analysis was locked to reward delivery (after the saccade). Analysis was segregated on the basis of the features that OFC neurons seemed most sensitive to: choice, whether that choice repeated the last choice and whether reward was delivered. For example, one of the conditions corresponded to rewarded saccades to a red target when that target was chosen last time. For the third and fourth MT datasets, stimuli were random dot patterns. For MT direction/area, stimulus direction and dot-patch area varied across conditions. For the MT speed dataset, stimuli were presented at different speeds (1–128 degrees per s) in the preferred and null directions. For these two datasets, the exact location of the individual dots in the aperture was different on each trial. This presumably did inflate the Fano factor by injecting additional across-trial variability. However, this appears to be a modest concern: robust Fano factor declines were still observed. These two datasets were also unusual in that the stimulus (the dot patch) appeared at the moment the trial began (and data collection commenced). Motion onset occurred 256 ms later.

Extracellular V1, MT direction/area and MT speed datasets were recorded from opioid-anesthetized monkeys (*macaca fascicularis* or *macaca nemestrina*). Intracellular V1 recordings were made from anesthetized cats. All others were made from awake monkeys (*macaca mulatta*) performing a task with controlled fixation. For the V4 dataset, the task was simply to fixate while stimuli were presented. For PMd, the task required a reach. For PRR, the task required a reach or a saccade. For the remaining datasets, the task required a saccade indicating a choice or perceptual discrimination. Tasks involved a delay between stimulus onset and when a response was allowed, with one exception. For the MT dots dataset, the monkey was allowed to respond as soon as he had made the discrimination. Analysis was thus limited to reaction times longer than 250 ms to allow a sufficient analysis window. All tasks involved a period of enforced fixation before stimulus onset. For LIP, this was not true for all trials and analysis was thus restricted to those that did. Fixation enforcement makes it unlikely that changes in variability are indirectly the results of saccades. In addition, datasets recorded from anesthetized/paralyzed animals showed the same changes in variability. Finally, it is unlikely that changes in the Fano factor, with a latency of ~60 ms, could be indirectly the results of saccades, which would incur a latency of 80–200 ms⁴⁹ plus an additional visual delay of 40–80 ms before firing rates could be affected. For all analyses, each trial's data were aligned on stimulus onset.

Most datasets were recorded across multiple days (and more than one monkey for LIP and MT) using single electrodes. Exceptions were the V1 and PMd datasets. The V1 data (dataset 565L) were recorded in 1 d from a 96-channel electrode array (Blackrock Microsystems). Only single-unit isolations (44) were analyzed. The PMd dataset was recorded using a similar array. Data were pooled across 5 d to increase trial and neuron count (although many units were probably recorded repeatedly). The single-neuron counts for the datasets used for the Fano factor analysis were 44 (V1), 49 (V4), 314 (MT plaids), 82 (MT dots), 43 (LIP), 175 (PRR), 73 (PMd), 116 (OFC), 47 (MT direction/area) and 57 (MT speed). The mean number

of trials per condition was 100 (V1), 24 (V4), 15 (MT plaids), 88 (MT dots), 35 (LIP), 10 (PRR), 31 (PMd), 106 (OFC), 125 (MT direction/area) and 14 (MT speed). The total numbers of neuron trials (number of neurons times the number of trials per neuron) were 52,800 (V1), 7,063 (V4), 106,007 (MT plaids), 7,179 (MT dots), 8,357 (LIP), 14,160 (PRR), 35,953 (PMd), 49,389 (OFC), 61,343 (MT direction/area) and 12,797 (MT speed).

Factor analysis requires simultaneously recorded data (for example, array-based recordings from a single day). Simulations similar to that in **Supplementary Figure 5** convinced us that the V1 and PMd datasets used for the Fano factor analyses had neuron and trial counts barely sufficient to provide factor analysis with sufficient statistical power. To increase statistical power, we included multi-unit isolations in the analysis. We used the two PMd recording days with the most (>~50) trials per condition (G20040123 and G20040122). For V1, we employed two datasets (567R and 106R). Both used the same stimuli as the dataset used for the Fano factor, but had higher unit and trial counts. The first (567R) had only 20 single-unit isolations, but many multi-unit isolations (102) and more trials per condition (200 rather than 100) relative to the dataset used for the Fano factor analysis. The second dataset (106R) had a very large (172) number of primarily multi-unit isolations, and even more trials per condition (400). These datasets were thus well-suited to factor analysis. Both of these datasets exhibited very clear Fano factor declines (**Supplementary Fig. 6**). However, the decision was made to keep the original Fano factor analysis (**Fig. 3**) restricted to the single unit-only dataset. Although the Fano factor can be reliably computed for multi-unit recordings (the sum of independent Poisson-distributed counts is Poisson distributed), the mean-matching procedure is not guaranteed to work.

Fano factor. The Fano factor was computed using Matlab (Mathworks) code available at (<http://www.stanford.edu/~shenoy/GroupCodePacks.htm>). Data for each neuron and condition are initially treated separately. A condition corresponds to multiple trials of the same stimulus. Spike counts were computed in a 50-ms sliding window moving in 10-ms steps. We then computed the variance (across trials) and mean of the spike count. Thus, the data for a particular neuron, condition and time were reduced to two numbers: the variance and mean of the spike count in a window centered on that time. For each time, a scatter plot of the variance versus the mean was compiled, with one point per neuron and condition (typically a few thousand points). The raw Fano factor was the slope of the regression relating the variance to the mean. This regression was weighted according to the estimated sampling error for the variance (computed on the basis of the number of trials and the mean rate, and assuming Poisson statistics). The observed effects did not result from the use of a short (50 ms) window; effect magnitudes were generally larger with larger windows, at the cost of blurred time courses.

To compute the mean-matched Fano factor, we first computed the distribution of mean counts (one datum per neuron and condition) for each time being analyzed (**Fig. 4b**). We then computed the greatest common distribution present at all times. Each bin of this common distribution had a height equal to the smallest value for that same bin across all distributions at all times. For each time, we then matched the analyzed distribution of mean rates to this common distribution. This was accomplished by discarding, from each bin, randomly chosen data points (each corresponding to the variance and mean pair for one neuron and condition) until the height of that bin matched that of the common distribution. The Fano factor was then computed. This procedure was repeated 50 times with different random seeds, with a different subset of the data preserved on each repetition, and thus a slightly different resulting Fano factor. The reported Fano factor is the mean of these individual Fano factors.

This mean-matching method discards less data than might be expected and far less than if we restrict analysis to individual neurons whose rates were not changing for particular conditions (as in **Fig. 5**). Mean matching preserved 49% (V1), 31% (V4), 71% (MT plaids), 32% (MT dots), 48% (LIP), 84% (PRR), 75% (PMd), 86% (OFC), 46% (MT-direction/area) and 70% (MT-speed) of the combinations of neuron and condition. Prior studies using the Fano factor have occasionally presented controls involving informal mean matching, drawing attention to different values of the Fano factor over a similar range of rates^{9,21,34}. Our method removes the need for such additional controls. Mean matching also does more than validate the basic effect; it prevents artifacts from corrupting the time course (**Supplementary Fig. 2**).

The statistical significance of the Fano factor decline was assessed by comparing the Fano factor 100 ms before stimulus onset and 200 ms after (for the MT

direction/area and MT speed datasets, where the Fano factor underwent two declines, we compared the time of motion onset with 200 ms after). Significance was computed on the basis of sampling distributions estimated from the 95% confidence intervals yielded by the regression used to compute the Fano factor. The decline was statistically significant in all cases: $P < 0.001$ (V1), $P < 10^{-5}$ (V4), $P < 10^{-10}$ (MT plaids), $P < 0.01$ (MT dots), $P < 0.005$ (LIP), $P < 0.002$ (PRR), $P < 10^{-10}$ (PMd), $P < 0.02$ (OFC), $P < 10^{-14}$ (MT direction/area) and $P < 10^{-7}$ (MT speed).

Factor analysis. Factor analysis was applied to spike counts in a 400-ms window, which either ended at stimulus onset (prestimulus) or began 100 ms after stimulus onset (stimulus). For the second V1 dataset (Fig. 6c), stimulus duration was only 400 ms, so we used a shorter (350 ms) window that began 50 ms after stimulus onset. A data matrix, D , was compiled, with D_{ij} being the spike count for the i^{th} trial and the j^{th} neuron. D included only neurons with rates ≥ 1 spike per s for both periods. Factor analysis was then applied to this matrix (neurons are variables, trials are observations). Factor analysis was performed separately on the D matrices compiled before and after stimulus onset. Factor analysis was performed separately for each condition. Subsequent results were averaged. We used previously published code²² to fit the parameters using expectation-maximization. Essentially identical results were obtained using Matlab's 'factoran'. Factor analysis requires an estimate of the dimensionality of the latent space capturing shared variability. To estimate this, we exploited the fact that recordings were made for multiple stimulus conditions (14 for PMd, 12 and 8 for the two V1 datasets). We computed the mean (across time and trials) response for each condition. We also included the pre-stimulus period as a condition. We then applied principal component analysis (to the $c \times n$ matrix) and asked how many dimensions were necessary to account for 95% of the across-condition response variance (principal component analysis is acceptable because it is being applied to trial-averaged data). For PMd and V1, five and four dimensions were required. These were then the latent dimensionalities assumed by factor analysis. These numbers likely underestimate the true dimensionality of the network response across conditions (our stimuli spanned only part of the space of all stimuli). This underestimate is acceptable, as it implies that factor analysis will underestimate the drop in the network variance, making our current interpretation conservative (Supplementary Fig. 7). Finally, similar results were obtained across a range of reasonable latent dimensionalities (2–8).

We used factor analysis to estimate the private and shared component of each neuron's variance (that is, we computed, but did not analyze, the full network covariance matrix). This kept each variance associated with a mean rate, and allowed us to apply the mean-matching procedure used for the Fano factor. Mean matching aids interpretation because even when rates are rising markedly, we can ask whether rate variance is changing for a given mean rate. Furthermore, there is a clear expectation that the estimated private noise should change little (assuming effects are not due to private noise), and there is less concern that a large increase in private noise (which will occur with rising mean rates) might contaminate the network variance estimate. The percentage of neurons and conditions preserved following mean matching was 31% and 48% (two V1 datasets) and 74% and 79% (two PMd datasets).

The estimated variances plotted in Figure 6b,c,e,f are the mean network (shared) variances, averaged across neurons and conditions, after mean matching. Significance was tested via t test (n = number of conditions).

GPFA. GPFA²⁶ was used to extract temporally smooth single-trial neural trajectories. Spike counts were taken in non-overlapping 20-ms bins, then square-rooted. The square-root transform⁵⁰ allows GPFA (which assumes stationary private noise for each neuron) to handle what would otherwise be a time-varying scaling of private noise with changing rates. Unlike factor analysis, GPFA was applied simultaneously to data from all time points. It was critical that factor analysis was applied separately before and after stimulus onset, to allow it to capture any changes in private noise. For GPFA, we require a visualization in which the same latent space captures the neural trajectory across time. Using all time points also increases the quantity of analyzable data. The cost is that GPFA, unlike the factor analysis analysis, cannot account for changes in private noise with time. Such changes are not the focus of the GPFA analysis (which seeks to track changes in network state, not private noise).

Figures 7 and 8 plot two-dimensional projections of the higher-dimensional trajectories found by GPFA. For PMd, subsequent analysis quantified ellipsoid volumes at different times. This analysis employed the higher-dimensional latent space. The dimensionality of that space was estimated using cross-validation. This is reasonable for GPFA, relative to factor analysis (where we did not use cross-validation), because of the increase in data points gained by including different times. Across the 14 reach targets, each analyzed separately, the latent dimensionality ranged from 8 to 12. It is expected that this dimensionality is higher than that for factor analysis, as the latter needed only to account for variability at a single time, while GPFA must account for variability across both trials and time. GPFA produced similar results when we used a dimensionality smaller (6) or higher (15) than that chosen by cross validation. Because dimensionality affects the covariance volume, we report that volume per dimension (i.e., volume^{1/k} where k is the latent dimensionality; k was always the same pre- and post-target).

For PMd, the two-dimensional projections in Figure 7 were chosen to preserve the relative per-dimensional volumes of the 8–12 dimensional covariance ellipses. For V1, we simply plotted the top two dimensions returned by GPFA, which maximizes the captured variability (analogous to the first two principal components).

43. Sugrue, L.P., Corrado, G.S. & Newsome, W.T. Matching behavior and the representation of value in the parietal cortex. *Science* **304**, 1782–1787 (2004).
44. Cohen, M.R. & Newsome, W.T. Context-dependent changes in functional circuitry in visual area MT. *Neuron* **60**, 161–173 (2008).
45. Armstrong, K.M., Fitzgerald, J.K. & Moore, T. Changes in visual receptive fields with microstimulation of frontal cortex. *Neuron* **50**, 791–798 (2006).
46. Armstrong, K.M. & Moore, T. Rapid enhancement of visual cortical response discriminability by microstimulation of the frontal eye field. *Proc. Natl. Acad. Sci. USA* **104**, 9499–9504 (2007).
47. Chang, S.W., Dickinson, A.R. & Snyder, L.H. Limb-specific representation for reaching in the posterior parietal cortex. *J. Neurosci.* **28**, 6128–6140 (2008).
48. Priebe, N.J., Churchland, M.M. & Lisberger, S.G. Constraints on the source of short-term motion adaptation in macaque area MT. I. The role of input and intrinsic mechanisms. *J. Neurophysiol.* **88**, 354–369 (2002).
49. Boch, R. & Fischer, B. Saccadic reaction times and activation of the prelunate cortex: parallel observations in trained rhesus monkeys. *Exp. Brain Res.* **50**, 201–210 (1983).
50. Kihlberg, J.K., Herson, J.H. & Schotz, W.E. Square root transformation revisited. *Appl. Stat.* **21**, 76–81 (1972).